

Respiratory Syncytial Virus (RSV) Guidance for Childcares

Respiratory Syncytial Virus (RSV) is a viral disease that primarily

affects children but can occur in people of all ages. Most people recover in a week or two, but RSV can be serious, especially for infants and older adults. Most children have been infected with RSV by age two and usually during the winter and early spring months (November-April).

Important Considerations	Prevention of RSV		
 Children often are infectious before symptoms appear. Excluding children with colds or other respiratory illnesses will probably NOT decrease the likelihood of transmission. You should exclude until the child has no fever and can tolerate normal activities. A child does <u>not</u> have to have a negative RSV test to return to childcare. 	 Careful and frequent hand washing! (warm water, soap, wash hands for 20 seconds) Proper disposal of tissues used to clean the nose and respiratory secretions. Prevent children from sharing cups and eating utensils with others. Clean and disinfect all surfaces on a regular basis, as well as after contact with ill children. Clean and disinfect toys on a regular basis, as well as after contact with ill children. 		

Follow the guidance in the <u>School and Childcare Exclusion List</u> for when children should be excluded and can return to childcare.

Disease/ Agent	Mode of Transmission	Contagious Period	Report to Public Health	Exclusion	Documentation for Return
RSV (Respiratory Syncytial Virus)	Respiratory (droplet) route or contact with infected individuals or contaminated objects	Shed for 3-8 days for children and adults May shed for 3-4 weeks in young infants and in immuno- suppressed individuals	Report outbreaks IMMEDIATELY by phone	Exclude younger children with RSV if the child has a <u>fever</u> or if the child is too sick to participate in activities with other children and staff.	Parent note

Contact the DHEC Health Department that is local to your area if you have additional questions.

<u>Lowcountry</u>	<u>Midlands</u>	<u>Pee Dee</u>	Upstate
Allendale, Bamberg, Beaufort,	Aiken, Barnwell, Chester,	Clarendon, Chesterfield,	Abbeville, Anderson, Cherokee,
Berkeley, Calhoun, Charleston,	Edgefield, Fairfield, Lancaster,	Darlington, Dillon, Florence,	Greenville, Greenwood, Laurens,
Colleton, Dorchester, Hampton,	Lexington, Kershaw, Newberry,	Georgetown, Horry, Lee, Marion,	McCormick, Oconee, Pickens,
Jasper, Orangeburg	Richland, Saluda, York	Marlboro, Sumter, Williamsburg	Spartanburg, Union
4050 Bridge View Drive, Suite 600	2000 Hampton Street	1931 Industrial Park Road	352 Halton Rd.
N. Charleston, SC 29405	Columbia, SC 29204	Conway, SC 29526	Greenville, S.C. 29607
Office: (843) 441-1091	Office: (888) 801-1046	Office: (843) 915-8886	Office (864) 372-3133
Fax: (843) 953-0051	Fax: (803) 576-2993	Fax: (843) 915-6506	Fax: (864) 282-4373
Nights/Weekends: (843) 441-1091	Nights/Weekends: (888) 801-1046	Nights/Weekends: (843) 409-0695	Nights/Weekends: (864) 423-6648

Reviewed 1/18/2023

S.C. Department of Health and Environmental Control